



# BEC Feed Solutions Pty Ltd.

50 & 66 Antimony Street. Carole Park Qld 4300

Ph: 07 3723 9800 Fax: 07 3271 3080

Email: [info@becfeedsolutions.com.au](mailto:info@becfeedsolutions.com.au)

Website: [www.becfeedsolutions.com.au](http://www.becfeedsolutions.com.au)

## GROMYCIN™

### Indications

For use in diets of Beef, Broilers and Dairy. For Cattle use to reduce lactic acidosis when high grain diets are fed.

### Composition

20 g / kg Virginiamycin

### Directions

#### Broilers and Turkeys

Day old to slaughter: Thoroughly mix 1.0kg of GROMYCIN™ premix to each 1 tonne of finished feed.

#### Grower Pigs

For pigs up to 34 kg liveweight: Thoroughly mix 1 kg of GROMYCIN™ premix to each 1 tonne of finished feed.

#### Cattle

Thoroughly mix 1 kg of GROMYCIN™ premix to each 1 tonne of finished feed.

For best results GROMYCIN™ premix should be thoroughly mixed with 10 kg of finely ground millrun or grain and then incorporated into the finished feed. Use mixed feed within 21 days of mixing.

### Discussion

When used in Broilers and Turkeys it has been shown that GROMYCIN™ increases weight gain and improves feed conversion. This is achieved by suppressing growth inhibiting bacteria and allowing better

absorption of nutrients. GROMYCIN™ will also slow down the transit time of foodstuffs through the gut.

When GROMYCIN™ is used in cattle feeds, it controls the growth of lactate producing bacteria in the rumen and caecum. This process leads to a reduction in lactic acidosis in cattle, especially when fed high carbohydrate grains. GROMYCIN™ is compatible with other feed additives in finished feeds.

### Withholding Period

Meat - Nil  
Eggs - Do not feed to birds laying eggs for human consumption.

### Schedule

S4

### Storage

Store below 30°C (Room Temperature). Protect from moisture.

### Expiry after Manufacture

12 months

### Pack Sizes

1 x 25 kg

